




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

Assessing the Collateral Damage of the Novel Coronavirus: A Call to Action for the Post-COVID-19 Era

Dominique Vervoort, MD, Jessica G.Y. Luc, MD, Edward Percy, MD, Sameer Hirji, MD, MPH, Richard Lee, MD, MBA



PII: S0003-4975(20)30618-4

DOI: <https://doi.org/10.1016/j.athoracsur.2020.04.015>

Reference: ATS 33740

To appear in: *The Annals of Thoracic Surgery*

Received Date: 21 April 2020

Accepted Date: 27 April 2020

Please cite this article as: Vervoort D, Luc JGY, Percy E, Hirji S, Lee R, Assessing the Collateral Damage of the Novel Coronavirus: A Call to Action for the Post-COVID-19 Era, *The Annals of Thoracic Surgery* (2020), doi: <https://doi.org/10.1016/j.athoracsur.2020.04.015>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 by The Society of Thoracic Surgeons

# **Assessing the Collateral Damage of the Novel Coronavirus: A Call to Action for the Post-COVID-19 Era**

Running Head: The collateral damage of COVID-19

Dominique Vervoort, MD<sup>1</sup>; Jessica G.Y. Luc, MD<sup>2</sup>; Edward Percy, MD<sup>2,3</sup>;  
Sameer Hirji, MD, MPH<sup>3</sup>; Richard Lee, MD, MBA<sup>4</sup>

<sup>1</sup> John Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

<sup>2</sup> Division of Cardiovascular Surgery, Department of Surgery, University of British Columbia, Vancouver, British Columbia, Canada

<sup>3</sup> Division of Cardiac Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

<sup>4</sup> Division of Cardiothoracic Surgery, Medical College of Georgia at Augusta University, Augusta, Georgia, USA

## **Corresponding Author:**

Dominique Vervoort, MD  
Johns Hopkins Bloomberg School of Public Health  
615 North Wolfe Street, 21205 Baltimore, Maryland  
[vervoortdominique@hotmail.com](mailto:vervoortdominique@hotmail.com)

**Word Count:** 2,307/2,500

**Keywords:** Cardiac Surgery; COVID-19; Coronavirus; Education

As of April 20th, 2020, the coronavirus disease 2019 (COVID-19), caused by SARS-CoV-2, has killed over 170,000 people worldwide and mortality projections for the United States have been as high as 200,000.<sup>1</sup> Health systems have been pressed with an uncomfortable reality to rapidly adapt to the growing burden of COVID-19 patients requiring critical care. This comes at the expense of elective surgeries, which are increasingly being cancelled across the world. This paper explores the contemporary issues impacting healthcare resources during the pandemic and makes recommendations for identifying an appropriate balance of their use to optimize patient care.

### Impact of Current Resource Allocation

Whilst necessary precautions must be taken to optimize patient care and the paucity of available resources in equipment, personnel, and infrastructure, collateral damage is inevitable (**Figure 1**). The following is anticipated in different time-frames or waves:

#### *First Wave: COVID-19 Infections and Burden on the Healthcare System*

- *CVD as a poor prognostic factor.* CVD is associated with a higher death rate in patients with COVID-19 (13.2%) compared to other comorbidities, including diabetes (9.2%), chronic respiratory disease (8.0%), hypertension (8.4%), and cancer (7.6%).<sup>2</sup> As such, the ethical obligation of physicians to prioritize the well-being of individual patients may be overridden by a delicate balance of policies that aim to prioritize the greatest good for the greatest number of patients: premature withdrawal or cessation of care in cardiovascular patients or 'Do not resuscitate'/'Do not intubate' in elderly with COVID-19 and perceived limited life expectancy, in order to reallocate resources to those with a better prognosis.<sup>7</sup> Conversely, by delaying patients with underlying CVD, they may be faced with a worse prognosis if they become infected with COVID-19.
- *A reduction in healthcare personnel, limiting access to care.* Healthcare workers themselves are at substantial risk for SARS-CoV-2 infection due to

increased exposure to high viral loads.<sup>2</sup> Additionally, increased rates of burnout among health workers, and cases of post-traumatic stress disorder as a result of frontline experiences can lead to a shortage of professionals trained to care for patients.<sup>3</sup>

- *Blood bank shortages.* Social distancing and lack of COVID-19 test kits for blood donors has led to a substantial decline in the availability of blood products. Although elective procedures requiring blood products are cancelled at most institutions, surgical emergencies may still necessitate blood reserves to optimize outcomes.

*Second Wave: Delayed Care and Presentation for Patients with CVD*

- *Delayed routine screening.* Recommended screening for malignancies (e.g., colorectal cancer), predisposed conditions (e.g., diabetes), and preexisting comorbidities (e.g., cardiac stress tests) may be partially postponed for an indefinite time until hospitals can reasonably manage existing patient loads. This will likely negatively impact their prognosis and lifespan.
- *Delayed presentations of patients with progressive disease.* For example, the recommendations for 'lockdown' to 'flatten the curve' may deter some patients (e.g., patients with chest pain, node positive lung cancer) from seeking time-sensitive care, out of fear or lack of access to transportation. They may be forced to or otherwise elect to "wait out their symptoms", which could pose substantial risk.
- *Declines in transplant procedures.* Given the risks associated with procurement team travel, combined with increased infection risk in recipients who will be immunosuppressed post-operatively, non-urgent transplantations may be delayed.

*Third Wave: Cardiothoracic Patients Awaiting Surgery or Losing Continuity of Care*

- *Reduction in preventive medicine.* The acute measures to stay at home and social distancing may lead to a myriad of negative effects in mental health,

physical wellbeing, and other lifestyle behaviours. Additionally, the shift in priority of primary care to triage suspected COVID-19 patients may limit the time available for routine visits (e.g., tracking CVD risk factors, such as hypertension, cholesterol, and diabetes – which have previously demonstrated survival benefit), supporting smoking cessation, and providing psychosocial support, despite the emergence of telehealth visits.

*Fourth Wave: Long-Term Post-COVID-19 Impact and Beyond*

- *Education and training.* As cardiothoracic surgical cases are cancelled and trainees are redeployed to areas of need, surgical trainees will increasingly find themselves with the challenge of balancing service and education. Some institutions are alternating residents and changing rotational structure. Coupled with a dramatic reduction in elective case volume, training opportunities are greatly diminished. Moreover, large and intermediate group didactic and process improvement meetings are routinely being cancelled. Since training in cardiothoracic surgery is over such a short amount of time—ranging from a minimum of two years via fellowship to a minimum of six years via integrated programs--the long-term impact of multiple lost months may be substantial.

As healthcare workers are redeployed to care for COVID-19 patients, and resources such as ventilators and extracorporeal membrane oxygenation devices are utilized to stabilize critically-ill COVID-19 patients, the capacity to treat CVD and emergencies remains critically limited. The decision to subject cardiovascular patients to surgery is difficult yet fluid, and with limited evidence-based guidelines as illustrated by the spectrum of CVD. While hemodynamically unstable patients are quickly moved to the operating room if pharmacological and non-invasive care has failed, the timing to intervene on clinically stable patients may not be as clear. In the context of heart transplants, United Network for Organ Sharing status 1 to 3 patients ought to be prioritized wherever possible, whilst those of lower

status are left in the dark.<sup>4</sup> Newborns with single ventricles or prostaglandin-dependence should clearly be managed, but what about children with valve insufficiency or large septal defects? Healthcare professionals are increasingly faced with important ethical dilemmas that are unique to the traditionally high-resource settings they have been operating in. The Society of Thoracic Surgeons has provided important ethical guidelines and tiered triage guidance statements for cardiac surgery, congenital heart surgery, and thoracic malignancies.<sup>3,5-7</sup> Furthermore, they have also developed important resource utilization prediction tools based on data from the Society of Thoracic Surgeons Database to assist healthcare providers in making informed decisions regarding resource allocation and triage of patients.

## Recommendations

As we move forward in this physical and emotional war, we propose the following for all healthcare professionals alike:

1. *Real-time outcomes tracking* will better inform the development of disease-specific guidelines for specialty care during pandemics, as well as the interaction between COVID-19 and pre-existing surgical conditions. Rightfully so, there is focused attention on outcomes tracking of patients who develop COVID-19. However, we urge the medical community that the tracking of outcomes of non-COVID-19 patients is equally important to better inform us of not only the mortality rates of COVID-19 but also its collateral damage because of patients' reluctance to seek care, or the absence of otherwise routine care. The use of digital-health technologies should be employed to longitudinally track patients initially scheduled for elective surgery. For example, the inclusion of real-time tracking for cardiac mortality would result in a real-time impact of decision making: i.e., if mortality from CVD dramatically increases, a corrective course may be developed in real-time.
2. *We must remain committed to, and available for, all patients--COVID-19 or not—as a way of ensuring population health.* This strategy may also assist in keeping

comorbidities under control, especially as their propensity complicates the potential COVID-19 disease severity course. Decentralization of care and the use of telemedicine to consult patients from their own homes is necessary to maintain an optimal patient-doctor relationship, whilst providing preventive care, triaging of new conditions, and comprehensive health counseling. These aspects may be supplemented by the use of emerging telemonitoring and cardiac-wearables which can also track patients' vital signs as well as serve as portable point-of-care testing. Additionally, dedicated multidisciplinary virtual triage rounds, if feasible, may be initiated to convene healthcare professionals to discuss the care of patients whose procedures have been cancelled or whose health status has changed.

3. *Patient and community education is critical* to flatten the epidemiological curve, whilst ensuring that patients understand when to seek care, led by example by physician leaders. Establishing virtual emergency care triage is essential, and can allow for timely patient evaluation without subjecting them to unnecessary travel to the emergency department. Empowering patients to self-access information via their electronic medical record and digitization of patient support groups via chat groups, smartphone applications, and real-time video-conferencing, and remote rehabilitation can help to improve patient engagement and health literacy.
4. *Hospitals have instituted patient-centered policies and should further develop guidelines* to centralize and streamline urgent cardiovascular care in this rapidly changing era. These include ensuring that a separate emergency team remains active and able to accept cases, minimizing crossover infections among patients and health workers, dedicating select hospitals, catheterization labs, operating rooms, and wards to non-COVID-19 patients, and allowing other healthcare personnel to become available when COVID-19 patients surge. Furthermore, it may include the need to have dedicated hospitals caring for COVID-19 patients only, whilst other hospitals outside heavily-hit areas take care of non-COVID-19 patients. Lastly,



attention should be paid to preparing patients for post-acute care and rehabilitation to reduce care fragmentation and offload the burden from acute-care centers.<sup>8</sup>

5. *A balance should be struck between cancelation and delaying of elective surgical procedures, and the reallocation of surgical patients to non-guideline-directed therapies (e.g., thrombolytics versus revascularization for acute coronary syndromes).* Likewise, early-stage malignancies, small aneurysms, and asymptomatic congenital heart defects may be postponed to prioritize urgent interventions and COVID-19 critical care. However, this must be weighed against timely intervention for patients with potentially life-threatening illnesses. The importance of the multidisciplinary heart-team approach, in close collaboration with the hospital ethics and palliative care committee, cannot be overstated.
6. *Professional societies need to continue to rapidly develop policies benefiting frontline workers to take appropriate action and effectively communicate them.* Initial examples have been set by the Cardiothoracic Surgery Network (CTSNet) and Canadian Cardiovascular Society—developing guidelines and townhalls--and the Society of Thoracic Surgeons—launching its Resource Utilization Tool for Cardiac Surgery and virtual COVID-19 resource portal—and can be supplemented by clear guidelines such as the joint consensus statement on thoracic malignancies from the Thoracic Surgery Outcomes Research Network, The Society of Thoracic Surgeons, and the American Association for Thoracic Surgery.<sup>7,9</sup> Comprehensive consensus statements and guidelines that provide further guidance in the management of all cardiothoracic surgical diseases in each stage of the surge of COVID-19 are needed and can help alleviate the COVID-19 burden wherever possible.
7. *The integrity of surgical training and the sustainability of the workforce should be maintained.* Trainee wellness and safety should be priority. In addition, restructuring of workflow and adaptation of educational milestones can help minimize disruption to surgical training.<sup>10</sup> E-technology can help provide continuity in academic and simulation curricula as well as allow flexibility in interviews and licensing

examinations. However, such virtual education platforms will require rapid adoption by programs and growing understanding and familiarity by both trainees and faculty. Furthermore, it is important to recognize that competency is not determined by case numbers.<sup>10</sup> Maintaining a rigorous standard for surgical competency during a pandemic calls for a thoughtful and coordinated national approach. This may highlight the need for development and integration of advanced simulation in present day training.

As the COVID-19 pandemic continues, clinicians and policymakers must be weary of the potential collateral damage on non-COVID-19 patients, health workers, and health systems. The decisions we make now will have important longitudinal implications on patient care as well as healthcare worker wellbeing. The surgical workforce should continue to take the necessary steps to protect oneself, COVID-19 patients, and, importantly, also non-COVID-19 patients. Cardiovascular surgical emergencies, traumatic injuries, obstetric complications, and congenital malformations will continue to occur, and will likely expose the weaknesses of the already burdened health system. However, sound judgement and calculated responses--proactive rather than reactive--are vital to ensure the robustness of our health system and to promote population health. We providers have the responsibility to all our patients to make sure our resource allocation does more good than harm.

**References:**

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis.* March 2020. doi:10.1016/S1473-3099(20)30120-1
2. World Health Organization. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19).; 2020. <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf>.
3. Drake D, Morrow CD, Kinlaw K, De Bonis M, Zangrillo A, Sade RM. Cardiothoracic Surgeons in Pandemics: Ethical Considerations. *Ann Thorac Surg.* April 2020. doi:10.1016/j.athoracsur.2020.03.006
4. The International Society for Heart & Lung Transplantation. Guidance for Cardiothoracic Transplant and Ventricular Assist Device Centers Regarding the SARS CoV-2 Pandemic.; 2020. [https://ishlt.org/ishlt/media/documents/SARS-CoV-2\\_Guidance-for-Cardiothoracic-Transplant-and-VAD-centers.pdf](https://ishlt.org/ishlt/media/documents/SARS-CoV-2_Guidance-for-Cardiothoracic-Transplant-and-VAD-centers.pdf).
5. Haft JW, Atluri P, Alawadi G, et al. Adult cardiac surgery during the COVID-19 Pandemic: A Tiered Patient Triage Guidance Statement. *Ann Thorac Surg.* April 2020. doi:10.1016/j.athoracsur.2020.04.003
6. Stephens EH, Dearani JA, Guleserian KJ, et al. COVID-19: Crisis Management in Congenital Heart Surgery. *Ann Thorac Surg.* April 2020. doi:10.1016/j.athoracsur.2020.04.001
7. COVID-19 Guidance for Triage of Operations for Thoracic Malignancies: A Consensus Statement from Thoracic Surgery Outcomes Research Network. *Ann Thorac Surg.* April 2020. doi:10.1016/j.athoracsur.2020.03.005
8. Percy ED,\* Luc JG,\* Vervoort D, Hirji S, Ruel M, Coutinho T. Post-Discharge Cardiac Care in the Era of COVID-19: How Should We Prepare? [In Press: *Canadian Journal of Cardiology*, April 2020].
9. Vervoort D, Ma X, Luc JGY, Zieroth S. Rapid Scholarly Dissemination and Cardiovascular Community Engagement to Combat the Infodemic of the COVID-19 Pandemic. *Can J Cardiol.* April 2020. doi:10.1016/j.cjca.2020.03.042

10. Nissen AP, Smith JA, Schmitto JD, et al. Global Perspectives on Cardiothoracic, Cardiovascular, and Cardiac Surgical Training. J Thorac Cardiovasc Surg. February 2020. doi:10.1016/j.jtcvs.2019.12.111

Journal Pre-proof

**Figure Legend**

**Figure 1.** The health impact due to the novel coronavirus disease 2019 (COVID-19) and collateral damage due to impeded access to non-COVID-19 care. The first wave represents COVID-19 patients during the initial and delayed COVID-19 phase. The second wave represents non-COVID-19 patients requiring urgent or emergent cardiothoracic surgical care, and patients with late presentations. The third wave represents stable patient cohorts awaiting surgery but losing continuity of care. The fourth wave represents the socioeconomic and mental health impacts on patients and healthcare workers, as well as implications for reduced clinical workload for cardiothoracic surgery trainees and fellows. Figure adapted with permission from Dr. Victor Tseng

(<https://twitter.com/vectorsting/status/1244671755781898241>).




Figure adapted with permission from Dr. Victor Tseng (<https://twitter.com/VectorStina/status/1244671755781898241>)